

AD 653104

LIKELIHOOD RATIO TESTS FOR
RESTRICTED FAMILIES

by

Richard E. Barlow

ORC 67-16
APRIL 1967

OPERATIONS RESEARCH CENTER

COLLEGE OF ENGINEERING

STATEMENT NO. 1

DISTRIBUTION OF THIS Document is Unlimited

ARCHIVE COPY

UNIVERSITY OF CALIFORNIA - BERKELEY

DDC
RECEIVED
JUN 16 1967
RECEIVED
C

LIKELIHOOD RATIO TESTS FOR RESTRICTED FAMILIES

by

Richard E. Barlow
Operations Research Center
University of California, Berkeley

April 1967

ORC 67-16

This research has been partially supported by the Office of Naval Research Contract Nonr-3656(18) with the University of California. Reproduction in whole or in part is permitted for any purpose of the United States Government.

ACKNOWLEDGMENT

We would like to gratefully acknowledge the help and advice of Gordon Crawford, Albert W. Marshall and David W. Walkup. Leonard Jacobson machine computed the percentage points in Tables 1 and 2 and performed the robustness study.

ABSTRACT

Likelihood ratio tests for the problem: $H_0: F \in \mathcal{F}_1$ versus $H_1: F \in \mathcal{F} - \mathcal{F}_1$ are defined for certain nonparametric families of distributions \mathcal{F} and \mathcal{F}_1 . In particular the likelihood ratio test is defined and shown to be unbiased when \mathcal{F}_1 denotes the exponential distributions (possibly truncated) and \mathcal{F} denotes the distributions with increasing failure rate. Comparisons are made with competing tests. The problem of testing for increasing failure rate average is also examined.

LIKELIHOOD RATIO TESTS FOR RESTRICTED FAMILIES

1. Introduction

Tests for composite hypotheses having optimal properties for finite samples have been obtained for various special problems by an important principle due to Neyman and Pearson (1928, 1933) called the likelihood ratio principle. This principle leads to the likelihood ratio test. Asymptotic properties of this test for parametric families of distributions can be found in Wilks (1962). Recently a conditional likelihood ratio test has been proposed for testing for trend in a stochastic process of Poisson type [Boswell (1966)]. This is a departure from the standard literature in that the underlying family of distributions considered is essentially nonparametric. The main result obtained is the asymptotic distribution of the likelihood ratio under the null hypothesis of no trend.

We consider likelihood ratio tests for certain geometrically restricted families of distributions. For example, let

$$\mathcal{F}_0 = \left\{ F \mid F(0) = 0 \text{ and } \frac{-\log [1-F(x)]}{x} \text{ nondecreasing in } x \geq 0 \right\}.$$

Then \mathcal{F}_0 is known as the IFRA (for increasing failure rate average) family of distributions. These distributions play an important role in the mathematical theory of reliability [Birnbaum, Esary, and Marshall (1966)]. However, not only is the family nonparametric but there is no sigma-finite measure relative to which all $F \in \mathcal{F}_0$ are absolutely continuous. Hence, the usual concept of maximum likelihood estimate does not suffice. Kiefer and Wolfowitz (1956, p. 893) propose a generalization of the maximum likelihood estimate concept which we adopt. Let $F_1, F_2 \in \mathcal{F}$ and let $f(\cdot; F_1, F_2)$ denote

the Radon-Nikodym derivative of F_1 with respect to the measure induced by $F_1 + F_2$.

Definition 1

\hat{F} is called the maximum likelihood estimate relative to \mathcal{F} if \hat{F} satisfies

$$\sup_{F \in \mathcal{F}} \prod_{i=1}^n \left[\frac{f(X_i; F, \hat{F})}{1 - f(X_i; F, \hat{F})} \right] = 1,$$

where $\underline{X} = (X_1, X_2, \dots, X_n)$ is a random sample from some $F \in \mathcal{F}$.

This definition is easily seen to coincide with the usual definition when the family \mathcal{F} is dominated by a sigma-finite measure.

Now consider the problem of testing $H_0: F \in \mathcal{F}_0$ against the alternative $H_1: F \in \mathcal{F} - \mathcal{F}_0$ where $\mathcal{F}_0 \subset \mathcal{F}$. Let $\hat{F}_0(\hat{F})$ denote the maximum likelihood estimate relative to \mathcal{F}_0 (\mathcal{F}) in the sense of definition 1. We define the likelihood ratio statistic $\Lambda_n(\underline{X})$ based on a random sample \underline{X} as follows:

Definition 2

$\Lambda_n(\underline{X})$ is called the likelihood ratio statistic where

$$\Lambda_n(\underline{X}) = \prod_{i=1}^n \left[\frac{f(X_i; \hat{F}_0, \hat{F})}{1 - f(X_i; \hat{F}_0, \hat{F})} \right].$$

We will be concerned with the properties of $\Lambda_n(\underline{X})$ for various restricted families of distributions $\mathcal{F}_0 \subset \mathcal{F}$.

2. IFRA Distributions

Let $\mathcal{F}_0 = \{F \mid F(0) = 0 \text{ and } \frac{-\log [1-F(x)]}{x} \uparrow \text{ in } x \geq 0\}$ and $\underline{X} = (X_1, X_2, \dots, X_n)$ denotes a random sample from F . We claim that the maximum likelihood

estimate (MLE) \hat{F}_0 relative to \mathcal{F}_0 puts mass at each of the sample observations. To see this suppose $\mathcal{F}^* = \{F_1, F_2\}$ where the F_1 probability of the observation X_i is $F_1\{X_i\} > 0$ for $i = 1, 2, \dots, n$ and $F_2\{X_k\} = 0$ for some k ($1 \leq k \leq n$). From definition 1 it follows that F_1 is MLE in \mathcal{F}^* . Since $F \in \mathcal{F}_0$ can put mass at a countable number of points we may restrict attention to those $F \in \mathcal{F}_0$ putting mass at sample points; i.e., F absolutely continuous with respect to $\lambda + \mu$ where λ is Lebesgue measure and $\mu\{A\}$ equals the number of sample points in A . The likelihood becomes

$$L_n(\underline{X} | F) = \prod_{i=1}^n F\{X_i\}.$$

Proschan and Marshall (1967) have obtained the MLE under the IFRA assumption. From the definition of IFRA distributions we see that

$$(2.1) \quad L_n(\underline{X} | F) = \prod_{i=1}^n \left[\exp(-\lambda_{i-1}X_i) - \exp(-\lambda_iX_i) \right]$$

where $0 \leq \lambda_0 \leq \lambda_1 \leq \dots \leq \lambda_n$. We maximize likelihood subject to these restrictions by letting $\lambda_0 = 0$ and $\lambda_n = +\infty$. Letting $\Delta\lambda_j = \lambda_j - \lambda_{j-1}$ where $\lambda_0 = 0$ and $\lambda_n = +\infty$, we see that (2.1) becomes

$$(2.2) \quad L_n(\underline{X} | F) = \prod_{i=1}^{n-1} \left[\exp(-\Delta\lambda_i \sum_{j=i+1}^n X_j) [1 - \exp(-\Delta\lambda_i X_i)] \right].$$

Maximizing (2.2) subject to $\Delta\lambda_i \geq 0$ ($1 \leq i \leq n$) we see that

$$(2.3) \quad \Delta \hat{\lambda}_i = \frac{1}{\hat{x}_i} \left[\log \sum_{j=1}^n x_j - \log \sum_{j=i+1}^n x_j \right] \geq 0$$

are the maximum likelihood estimates of $\Delta \lambda_i (1 \leq i \leq n)$. Substituting (2.3) in (2.2) we see that the maximum likelihood according to definition 1 becomes

$$(2.4) \quad L_n(\underline{X} \mid \hat{F}_1) = \prod_{i=1}^{n-1} \left[1 - \frac{x_i}{\sum_{j=i}^n x_j} \right]^{\frac{\sum_{j=i}^n x_j}{x_i} - 1} \left[\frac{x_i}{\sum_{j=i}^n x_j} \right].$$

If we let \mathcal{F} denote the class of all distributions on the positive axis, then \hat{F} is the usual empirical distribution function and the likelihood ratio statistic for testing H_0 versus H_1 becomes

$$(2.5) \quad \Lambda_n(\underline{X}) = n^n \prod_{i=1}^{n-1} \left[1 - \frac{x_i}{\sum_{j=i}^n x_j} \right]^{\frac{\sum_{j=i}^n x_j}{x_i} - 1} \left[\frac{x_i}{\sum_{j=i}^n x_j} \right].$$

We consider the test, ϕ , which rejects H_0 when

$$\Lambda_n(\underline{X}) \leq c_\alpha$$

where c_α is determined by

$$P_G \{ \Lambda_n(\underline{Y}) \leq c_\alpha \} = \alpha$$

and $G(x) = 1 - e^{-x}$ for $x \geq 0$. This choice of a test based on the likelihood ratio statistic is motivated by the following theorem.

Theorem 2.1

The test, ϕ , based on the likelihood ratio statistic $\Lambda_n(\underline{X})$ is unbiased at all significance levels for the problem

$$H_0 : F \in \mathcal{F}_0 = \{\text{IFRA}\}$$

versus

$$H_1 : F \in \mathcal{F}_1 = \{\text{DFRA}\}.$$

Proof:

Let $G(x) = 1 - e^{-x}$ for $x \geq 0$. It is sufficient to show that $F_0 \in \mathcal{F}_0$ implies

$$(2.6) \quad P_G \{ \Lambda_n(\underline{Y}) \leq x \} \geq P_{F_0} \{ \Lambda_n(\underline{X}) \leq x \}$$

and $F_1 \in \mathcal{F}_1$ implies

$$(2.7) \quad P_G \{ \Lambda_n(\underline{Y}) \leq x \} \leq P_{F_1} \{ \Lambda_n(\underline{X}) \leq x \}.$$

Let $\psi(x) = G^{-1}F_0(x)$, $Y_i^* = \psi(X_i)$ and note $\frac{\psi(x)}{x} \uparrow$ in $x \geq 0$. Also Y_i^* is distributed as the i -th order statistic from G . Now

$$\frac{\psi(X_i)}{X_i} \leq \frac{\psi(X_j)}{X_j} \quad \text{for } j \geq i$$

implies

$$\frac{\frac{x_i}{\sum_{j=i}^n x_j}}{\frac{\psi(x_i)}{\sum_{j=i}^n \psi(x_j)}} = \frac{\frac{y_i^*}{\sum_{j=i}^n y_j^*}}{\frac{\psi(y_i^*)}{\sum_{j=i}^n \psi(y_j^*)}} .$$

Since $h(x) = x(1-x)^{\frac{1}{x}-1}$ is increasing in x ($0 \leq x \leq 1$), it follows that

$$\prod_{i=1}^{n-1} h\left(\frac{x_i}{\sum_{j=i}^n x_j}\right) \geq \prod_{i=1}^{n-1} h\left(\frac{y_i}{\sum_{j=i}^n y_j}\right)$$

where $(y_1 \leq y_2 \leq \dots \leq y_n)$ is an independent ordered sample from G . (2.6)

follows immediately and the proof of (2.7) is similar. ||

We say that $F_1 \leq_* F_2$ (i.e., F_1 is starshaped with respect to F_2) if $\frac{F_2^{-1}F_1(x)}{x}$ is nondecreasing for $x \geq 0$. From the proof of theorem 2.1 it follows that $F_1 \leq_* F_2$ implies

$$P_{F_1} \{ \Lambda_n(\underline{X}) \leq c_\alpha \} \leq P_{F_2} \{ \Lambda_n(\underline{X}) \leq c_\alpha \} .$$

Hence the power of the likelihood ratio test is greater at F_2 than at F_1 when $F_1 \leq_* F_2$. Percentage points for $-\log \Lambda_n(\underline{X})$ are given in Table 1.

Of course there are many unbiased tests of the IFRA hypothesis. Marshall, Walkup and Wets (1966) have characterized the class of all such tests. These are just the tests based on functions $f(x_1, x_2, \dots, x_n)$ having the properties:

- 1) f is homogenous;
- 2) $\sum_{i=1}^j x_i \frac{\partial f(x_1, x_2, \dots, x_n)}{\partial x_i} \geq 0$ for $j = 1, 2, \dots, n-1$
and all $x_1 \geq x_2 \geq \dots \geq x_n \geq 0$.

The test associated with f would reject H_0 if

$$f(X_1, X_2, \dots, X_n) \geq c$$

where c is some suitable critical number and $X_1 \geq X_2 \geq \dots \geq X_n$ are the usual order statistics labelled in reverse order.

3. IFR Distributions

Let $\mathcal{F} = \{F \mid F(0) = 0 \text{ and } -\log [1 - F(x)] \text{ is convex for } x \geq 0\}$. This is the class of IFR (for increasing failure rate) distributions. Proschan and Pyke (1965) have proposed a test for constant versus increasing failure rate. Their test is based on a statistic considered by M. G. Kendall (1938) and H. B. Mann (1945) and is essentially a rank test for trend. Proschan and Pyke show that their test is unbiased, has good large sample properties, and is competitive in this sense with certain parametric tests when the unknown distribution lies within some specified parametric family. However, sampling experiments indicate that their test does not have good discriminating power for relatively small samples. This is perhaps to be expected since their test does not use all of the information in the sample.

M. Boswell (1966) studied a similar problem concerning Poisson type processes. His statistic based on a conditional maximum likelihood ratio test is essentially the same as the likelihood ratio statistic studied in this section. The main result in Boswell's paper is a derivation of the asymptotic distribution of his test statistic. In contrast, we concentrate on small sample results.

Since IFR distributions can have a jump at the right hand end of their interval of support it is clear from definition 1 that we need only consider

estimators absolutely continuous with respect to Lebesgue measure on $[0, X_n)$ with a jump at X_n (see Barlow and Proschan, (1965), p. 26). Hence

$$L_n(\underline{X} \mid F) = \left[\prod_{i=1}^{n-1} f(X_i) \right] F\{X_n\}$$

where f is the density of F on $[0, X_n)$. Since

$$1 - F(x) = \exp \left[- \int_0^x r(u) du \right] \quad \text{where } r(u) = f(u) / [1 - F(u)]$$

for $0 \leq u < X_n$, we may write

$$f(x) = r(x) e^{-\int_0^x r(u) du} \quad 0 \leq x < X_n$$

and

$$F\{X_n\} = e^{-\int_0^{X_n} r(u) du}.$$

Hence

$$(3.1) \quad \log L_n(\underline{X} \mid F) = \sum_{i=1}^{n-1} \log r(X_i) - \sum_{i=1}^n \int_0^{X_i} r(u) du.$$

The problem of maximizing (3.1) subject to $r(x)$ nondecreasing was solved by Grenander (1956) and independently by Marshall and Proschan (1965). They show that the problem can be reduced to maximizing

$$\sum_{i=1}^{n-1} \log r(X_i) - \sum_{i=1}^{n-1} (n-i) (X_{i+1} - X_i) r(X_i)$$

subject to $r(X_1) \leq r(X_2) \leq \dots \leq r(X_{n-1})$. The maximum likelihood estimates are

$$\hat{f}_n(x_i) = \min_{v \geq i+1} \max_{u \leq i} \left[\frac{(v-u)}{(n-u)(x_{u+1} - x_u) + \dots + (n-v+1)(x_v - x_{v-1})} \right]$$

for $i = 1, 2, \dots, n-1$. The maximum likelihood is

$$(3.2) \quad L_n(\underline{x} \mid \hat{f}) = \left[\prod_{i=1}^{n-1} \hat{f}(x_i) \right] e^{-(n-1)}.$$

The exponent on e can be easily verified using the definition of \hat{f} and observing that

$$\sum_{i=1}^n \int_0^{x_i} \hat{f}(u) du = \sum_{i=1}^{n-1} (n-i) \int_{x_i}^{x_{i+1}} \hat{f}(u) du.$$

Let $\mathcal{F}_0 = \{F \mid F(0) = 0, F(x) = 1 - e^{-\lambda x} \text{ for } x < T \text{ and } F(T) = 1, \lambda > 0, T > 0\}$

Then \mathcal{F}_0 denotes the class of exponential distributions with possible truncation on the right. Consider now the problem of testing $H_0: F \in \mathcal{F}_0$ versus

$H_1: F \in \mathcal{F} - \mathcal{F}_0$. The choice of H_0 was determined by the fact that the MLE's \hat{F}_0 and \hat{F} are both absolutely continuous with respect to Lebesgue measure in $[0, x_n)$ and place mass at x_n . The likelihood under H_0 will be

$$L_n(\underline{x} \mid F_0) = \left[\prod_{i=1}^{n-1} \lambda e^{-\lambda x_i} \right] e^{-\lambda x_n}$$

and the maximum likelihood will be

$$(3.3) \quad L_n(\underline{x} \mid \hat{F}_0) = \left(\frac{n-1}{\sum_{i=1}^{n-1} x_i} \right)^{n-1} e^{-n}.$$

According to definition 2, the likelihood ratio statistic for testing for truncated exponentiality versus IFR and not truncated exponentiality will be

$$(3.4) \quad \Lambda_n^*(\underline{x}) = \frac{(n-1)^{n-1}}{\left(\sum_{i=1}^n x_i\right)^{n-1} \prod_{i=1}^{n-1} \hat{f}(x_i)} .$$

If $\frac{1}{(n-1)(x_2 - x_1)} \leq \frac{1}{(n-2)(x_3 - x_2)} \leq \dots \leq \frac{1}{(x_n - x_{n-1})}$ so that

$$\hat{f}(x_i) = \frac{1}{(n-i)(x_{i+1} - x_i)} \quad i = 1, 2, \dots, n-1 ,$$

then (3.4) becomes

$$\Lambda_n^{**}(\underline{x}) = \left(\frac{n-1}{\sum_{i=1}^n x_i}\right)^{n-1} \prod_{i=1}^{n-1} (n-i)(x_{i+1} - x_i) .$$

As in section 2 we consider the test, φ^* , which rejects H_0 when

$$\Lambda_n^*(\underline{x}) \leq c_\alpha$$

where c_α is determined by

$$P_G \{ \Lambda_n^*(\underline{Y}) \leq c_\alpha \} = \alpha .$$

The asymptotic distribution of $\Lambda_n^*(\underline{Y})$ can be found in Boswell (1966, p. 1572).

A table of percentage points obtained using Monte Carlo methods is contained in Table 2.

4. Unbiasedness of the Likelihood Ratio Test for IFR

Like the Proschan-Pyke test, the likelihood ratio test has greater power under the alternative than under the null hypothesis. To show this we need to introduce some auxiliary results.

Given a sequence of nonnegative real numbers $\{z_i\}_{i=1}^n$, plot $\sum_{j=1}^i z_j$ versus i and interpolate linearly between $(0, 0)$, $(1, z_1)$, \dots , $(n, z_1 + \dots + z_n)$. Let $\bar{z}_1 \geq \bar{z}_2 \geq \dots \geq \bar{z}_n$ denote the slopes of the least concave majorant to this graph in successive intervals. This operation converts the original sequence into a nonincreasing sequence and will be useful later on. For convenience, call $\bar{z}_1 \geq \bar{z}_2 \geq \dots \geq \bar{z}_n$ the Brunkized sequence after D. Brunk (see D. Brunk et. al. (1955)). Note that $\bar{z}_1 \geq \bar{z}_2 \geq \dots \geq \bar{z}_n$ can also be obtained by successive averaging of the original sequence until it becomes nonincreasing.

We say that $H(z_1, z_2, \dots, z_n)$ is a Schur function if

$$(z_i - z_j) \left(\frac{\partial H}{\partial z_i} - \frac{\partial H}{\partial z_j} \right) \geq 0$$

for all vectors \underline{z} and $i, j = 1, 2, \dots, n$. This concept is needed in the following useful lemma.

Lemma 1

Let (z_1, z_2, \dots, z_n) and $(z'_1, z'_2, \dots, z'_n)$ denote two nonnegative sequences such that

$$\sum_{i=1}^r z_i \geq \sum_{i=1}^r z'_i \quad \text{for } r = 1, 2, \dots, n-1$$

and

$$\sum_{i=1}^n z_i = \sum_{i=1}^n z'_i.$$

Then the inequalities are preserved under Brunkization;

$$\text{i.e.,} \quad \sum_{i=1}^r \bar{z}_i \geq \sum_{i=1}^r \bar{z}_i^{\top} \quad (r = 1, 2, \dots, n-1)$$

(i)

$$\sum_{i=1}^n \bar{z}_i = \sum_{i=1}^n \bar{z}_i^{\top} .$$

If H is a Schur function then

$$(ii) \quad H(\bar{z}_1, \bar{z}_2, \dots, \bar{z}_n) \geq H(\bar{z}_1^{\top}, \bar{z}_2^{\top}, \dots, \bar{z}_n^{\top}) .$$

Proof:

(i) is obvious since the least concave majorant to the $\{z_i\}$ sequence lies above the least concave majorant to the $\{z_i^{\top}\}$ sequence.

Since (i) holds, $\bar{z}_1 \geq \bar{z}_2 \geq \dots \geq \bar{z}_n$; $\bar{z}_1^{\top} \geq \dots \geq \bar{z}_n^{\top}$ and $\sum_{i=1}^n \bar{z}_i = \sum_{i=1}^n \bar{z}_i^{\top}$

we have (ii) by the Schur, Ostrowski theorem (see Ostrowski (1952)). //

Theorem 4.1

If $G^{-1}F(x)$ is convex for $x \geq 0$, $G(0) = F(0) = 0$ and $\underline{X}(\underline{Y})$ denotes a random sample from $F(G)$, then

$$\Lambda_n^*(\underline{X}) \leq_{st} \Lambda_n^*(\underline{Y}) .$$

Remark

This proves the likelihood ratio test Λ_n^* is unbiased since if F is IFR and $G(x) = 1 - e^{-x}$ for $x \geq 0$, then $G^{-1}F$ is convex on $x \geq 0$.

Proof:

Let $Y_i^* = G^{-1}F(X_i)$ and note $Y_{i \text{ st}}^* = Y_i$.

Let

$$z_i = \frac{(n-i)(x_{i+1} - x_i)}{\sum_{l=1}^{n-1} (n-l)(x_{l+1} - x_l)} \quad \text{and}$$

$$z_i^* = \frac{(n-i)(y_{i+1}^* - y_i^*)}{\sum_{l=1}^{n-1} (n-l)(y_{l+1}^* - y_l^*)}.$$

Since $y_i^* = G^{-1}F(x_i)$ and $G^{-1}F$ is convex

$$\frac{(n-i)(y_{i+1}^* - y_i^*)}{(n-i)(x_{i+1} - x_i)}$$

is increasing in $i = 1, 2, \dots, n$. It follows from lemma 3.7 (i) of Barlow and Proschan (1966) that

$$\frac{\frac{\sum_{i=1}^r z_i}{\sum_{i=1}^r z_i}}{\frac{\sum_{i=1}^{n-1} z_i}{\sum_{i=1}^{n-1} z_i}} \leq \frac{\frac{\sum_{i=1}^{n-1} z_i}{\sum_{i=1}^{n-1} z_i}}{\frac{\sum_{i=1}^{n-1} z_i}{\sum_{i=1}^{n-1} z_i}} = 1$$

and hence $\frac{\sum_{i=1}^r z_i}{\sum_{i=1}^r z_i} \geq \frac{\sum_{i=1}^r z_i^*}{\sum_{i=1}^r z_i^*}$ for $r = 1, 2, \dots, n-1$. Let $\{\bar{z}_i\}$ and $\{\bar{z}_i^*\}$ denote the Brunkized estimates of $\{z_i\}$ and $\{z_i^*\}$ respectively. Let

$$-H(x_1, x_2, \dots, x_{n-1}) = \prod_{i=1}^{n-1} x_i$$

and note that H is a Schur function. Since $\{z_i\}$ and $\{z_i^*\}$ satisfy the hypotheses of lemma 1, it follows that

$$-H(\bar{z}_1, \bar{z}_2, \dots, \bar{z}_{n-1}) \leq -H(\bar{z}_1^*, \bar{z}_2^*, \dots, \bar{z}_{n-1}^*) .$$

Hence

$$(4.1) \quad \frac{1}{\left[\sum_{i=1}^{n-1} (n-i)(x_{i+1} - x_i) \right]^{n-1} \prod_{i=1}^{n-1} \hat{f}(x_i)} \leq \frac{1}{\left[\sum_{i=1}^{n-1} (n-i)(y_{i+1}^* - y_i^*) \right]^{n-1} \prod_{i=1}^{n-1} \hat{f}(y_i^*)} .$$

Since

$$\frac{\sum_{i=0}^r (n-i)(y_{i+1}^* - y_i^*)}{\sum_{i=0}^r (n-i)(x_{i+1} - x_i)} \geq \frac{\sum_{i=1}^n y_i^*}{\sum_{i=1}^n x_i}$$

for $(1 \leq r \leq n-1)$ by lemma 3.7 (i') of Barlow and Proschan (1966) it follows that

$$(4.2) \quad \frac{\sum_{i=1}^{n-1} (n-i)(x_{i+1} - x_i)}{\sum_{i=1}^n x_i} \leq \frac{\sum_{i=1}^{n-1} (n-i)(y_{i+1}^* - y_i^*)}{\sum_{i=1}^n y_i^*} .$$

(4.1) and (4.2) together imply

$$\Lambda_n^*(\underline{x}) \leq \Lambda_n^*(\underline{y}^*) .$$

The theorem follows from $(Y_1, Y_2, \dots, Y_n) \stackrel{st}{=} (Y_1^*, Y_2^*, \dots, Y_n^*)$. //

Marshall, Walkup and Wets (1966) have characterized the class of unbiased tests for constant failure rate versus nondecreasing failure rate. These are based on functions $h(x_1, x_2, \dots, x_n)$ satisfying the conditions

- i) h is homogeneous;
- ii)
$$\sum_{i=1}^j (x_i - x_{j+1}) \frac{\partial h(x_1, \dots, x_n)}{\partial x_i} \geq 0 \quad j = 1, 2, \dots, n-1$$

for all $x_1 > x_2 > \dots > x_n > 0$. The corresponding test consists of rejecting exponentiality if $h(X_1, X_2, \dots, X_n) \leq c$ where c is a suitable critical number and $X_1 \geq X_2 \geq \dots \geq X_n$ are the order statistics labelled in reverse order.

5. Distribution of the Maximum Likelihood Ratio Statistic Under the Exponential Assumption

From the computations in Boswell (1966) it is clear that the distribution of Λ_n^* , even under the null hypothesis, is exceedingly complicated. For this reason we have had to use Monte Carlo methods to obtain the percentage points tabulated in Table 2. However, the distribution of Λ_n^* under H_0 is quite smooth as we show in

Theorem 5.1

The likelihood ratio statistic Λ_n^* has a nonincreasing density on $(0, 1)$ under the exponential assumption.

Proof:

Let $0 = W_0 \leq W_1 \leq \dots \leq W_n$ denote an ordered sample from the uniform distribution on $(0, 1)$. Let

$$U_i = W_i - W_{i-1} \quad i = 1, 2, \dots, n-1.$$

Then the random vector $(U_1, U_2, \dots, U_{n-1})$ has joint density

$$h(u_1, u_2, \dots, u_{n-1}) = \begin{cases} (n-1)! & \text{for } u_i \geq 0 \\ & i = 1, 2, \dots, n-1 \\ & 0 \leq u_1 + u_2 + \dots + u_{n-1} \leq 1 \\ 0 & \text{otherwise.} \end{cases}$$

Let $(\bar{U}_1, \bar{U}_2, \dots, \bar{U}_{n-1})$ denote the modified vector $(U_1, U_2, \dots, U_{n-1})$ after Brunkization and subject to $\bar{U}_1 \geq \bar{U}_2 \geq \dots \geq \bar{U}_{n-1}$.

The likelihood ratio statistic

$$\Lambda_n^*(Y) = \frac{1}{\left(\frac{\sum_{i=1}^n Y_i}{n-1} \right)^{n-1} \prod_{i=1}^{n-1} \hat{r}(Y_i)}$$

is distributed as $(n-1)^{n-1} \prod_{i=1}^{n-1} \frac{n-1}{i}$ under the exponential assumption. Notationally it will be convenient to replace $n-1$ by n . Hence we need only prove that

$$P \left\{ \prod_{i=1}^n \bar{U}_i \leq z \right\}$$

is concave in $z \in (0, 1)$. Let I denote the usual indicator set function and observe that

$$P \left\{ \prod_{i=1}^n \bar{U}_i \leq z \right\} = n! \int \int \dots \int_{\substack{u_i \geq 0 \\ 0 \leq u_1 + \dots + u_n \leq 1}} I[\bar{u}_1 \dots \bar{u}_n \leq z] du_1 \dots du_n.$$

Integrating out on u_n we have

$$P\left\{\prod_{i=1}^n \bar{u}_i \leq z\right\} = n! \iint_{\substack{u_i \geq 0 \\ u_1 + \dots + u_{n-1} \leq 1}} \dots \int f(u_1, \dots, u_{n-2}; z) du_1 \dots du_{n-1}$$

where

$$\begin{aligned} f(u_1, \dots, u_{n-1}; z) &= \int_{0 \leq u_n \leq 1 - u_1 - \dots - u_{n-1}} [\bar{u}_1 \dots \bar{u}_n \leq z] du_n \\ &= \min [1 - u_1 - \dots - u_{n-1}, u_n(z)] \end{aligned}$$

and $u_n(z)$ is the solution of $z = \bar{u}_1 \dots \bar{u}_{n-1} \bar{u}_n$

for fixed u_1, u_2, \dots, u_{n-1} .

We claim that z is a strictly increasing convex function of u_n and, therefore, that u_n is a strictly increasing concave function of z . It follows that $f(u_1, \dots, u_{n-1}; z)$ is a concave function of z for fixed $(u_1, u_2, \dots, u_{n-1})$. Hence

$$P\left\{\prod_{i=1}^n \bar{u}_i \leq z\right\} = n! \iint_{\substack{u_i \geq 0 \\ u_1 + \dots + u_{n-1} \leq 1}} \dots \int f(u_1, \dots, u_{n-1}; z) du_1 \dots du_{n-1}$$

is a concave function of z .

To show $z = \bar{u}_1 \dots \bar{u}_{n-1} \bar{u}_n$ is a convex function of u_n , define $(\bar{u}_1^*, \dots, \bar{u}_{n-1}^*)$ to be the Brunk modification of $(u_1, u_2, \dots, u_{n-1})$ subject to $\bar{u}_1^* \geq \bar{u}_2^* \geq \dots \geq \bar{u}_{n-1}^*$. Clearly z is piecewise convex for u_n in the intervals $[0, \bar{u}_{n-2}^*], [\bar{u}_{n-2}^*, \bar{u}_{n-3}^*], \dots, [1 - \bar{u}_1^* - \dots - \bar{u}_{n-1}^*, \bar{u}_1^*]$. It is therefore sufficient to show that z has a continuous derivative in u_n . We show that the right and left hand derivatives at $u_n = \bar{u}_{n-1}^*$ are equal. For $u_n < \bar{u}_{n-1}^*$

$$\frac{dz}{du_n} = \bar{u}_1^* \dots \bar{u}_j^* \left(\frac{u_{j+1} + \dots + u_{n-1}}{n-1-j} \right)^{n-j-1}.$$

For $u_{n-1}^* < u_n < u_{n-2}^*$

$$\frac{dz}{du_n} = \bar{u}_1^* \dots \bar{u}_j^* \left(\frac{u_{j+1} + \dots + u_n}{n-j} \right)^{n-j-1}.$$

For $\bar{u}_n = \bar{u}_{n-1}^*$, obviously

$$\left(\frac{u_{j+1} + \dots + u_{n-1}}{n-1-j} \right) = \left(\frac{u_{j+1} + \dots + u_{n-1} + u_n}{n-j} \right). \quad ||$$

For $n = 2$ and $n = 3$ it is a straightforward computation to obtain the distribution of $\Lambda_n^*(Y)$. Clearly, for $n = 2$

$$\Lambda_2^*(Y) = U_1$$

and the likelihood ratio is uniformly distributed on $(0, 1)$.

For $n = 3$

$$\Lambda_3^*(Y) = 4 \bar{U}_1 \bar{U}_2 = \begin{cases} 4 U_1 U_2 & \text{if } U_1 \geq U_2 \\ (U_1 + U_2)^2 & \text{if } U_1 \leq U_2 \end{cases}.$$

Hence

$$P_G \left\{ \Lambda_3^*(Y) \leq u \right\} = \iint_{\left\{ \begin{array}{l} u_1 u_2 \leq \frac{u}{4} \\ u_1 \geq u_2 \end{array} \right\}} 2 u_1 \, du_1 \, du_2 + \iint_{\left\{ \begin{array}{l} u_1 + u_2 \leq \sqrt{u} \\ u_1 \leq u_2 \end{array} \right\}} 2 \, du_1 \, du_2$$

and

$$P_G \left\{ \Lambda_3^*(Y) \leq u \right\} = \frac{3u}{4} + \frac{1}{2} u \left[\log \left(\frac{1 + \sqrt{1-u}}{\sqrt{u}} \right) \right] + \left(\frac{1 - \sqrt{1-u}}{2} \right)^2.$$

The density is

$$g_3(u) = \frac{1}{2} + \frac{1}{2} \log \left[\frac{1 + \sqrt{1-u}}{u} \right] .$$

It is easy to check that g is decreasing, $g_3(0) = +\infty$, $g_3(1) = \frac{1}{2}$ and $g_3'(0) = g_3'(1) = -\infty$. It is tempting to conjecture that this behavior is true in general, i.e., $g_n(0) = +\infty$, $g_n'(0) = g_n'(1) = -\infty$ for $n \geq 3$.

TABLE 1 [†]
 Percentage Points
 for $-\log \Lambda_n(\underline{Y})$

Sample Size n	Percentiles			
	.01	.05	.90	.99
2	0.015	0.072	3.3	4.85
3	0.216	0.480	5.25	7.25
4	0.6	1.1	6.9	9.0
5	1.2	1.8	8.6	10.8
6	1.7	2.5	10.1	12.6
7	2.4	3.3	11.7	14.3
8	3.0	4.2	13.2	15.8
9	3.7	5.0	14.7	17.6
10	4.6	5.8	16.1	19.2

[†] Note that we use lower percentiles for testing exponentiality versus IFRA and upper percentiles for testing IFRA versus DFRA.

TABLE 2
 Percentage Points for the IFR
 Likelihood Ratio Statistic $\Lambda_n^*(\underline{y})$

Sample Size n	Percentiles		Number of Random Simulations Used
	.05	.01	
2	.0500	.0100	50,000
3	.025	.004	50,000
4	.0162	.0027	40,000
5	.0125	.0017	50,000
6	.01	.0015	60,000
7	.0087	.001	60,000
8	.0077	.001	80,000
9	.0065	.0007	70,000
10	.0055	.0007	50,000

6. Comparisons with Competing Tests

In a paper in process (Barlow and Jacobson) we study in some detail the robustness of the IFRA and IFR likelihood ratio tests relative to selected competing tests. Preliminary investigations indicate that the IFR likelihood ratio test is much better than the Proschan-Pyke test for small samples. However, they have achieved a remarkable amount of information concerning the asymptotic behavior of their test statistic. In particular they show that their test statistic, suitably normalized, has an asymptotic normal distribution for a wide class of alternative distributions. On the other hand, the distribution of the IFR likelihood ratio statistic, under the null hypothesis, converges to a weighted sum of χ^2 distributions which is rather cumbersome to compute. For small sample sizes (say $n \leq 10$) both statistics are relatively easy to compute with the aid of a desk calculator.

Figure 1 is a graph of the power functions for both tests against Weibull distribution alternatives (i.e., $F(x) = 1 - e^{-(\lambda x)^\alpha}$ for $x \geq 0$ where α is the shape parameter) when the sample size is 10 and the significance level is 5%. These curves were obtained by means of Monte Carlo simulation on a computer. The power of the Proschan-Pyke test increases very slowly as a function of the Weibull shape parameter, α . Even when the shape parameter $\alpha = 3$, the power is only .62. Our numerical investigations indicate that for $\alpha > 3$ the power increases even more slowly and that for $\alpha = 500$, the power is only .86. For $n = 20$, the power of the Proschan-Pyke test is much better; it yields a power of .92 for $\alpha = 3$. It should be noted, however, that the power of this test is still not as good as the power of the likelihood ratio test when $n = 10$.

The asymptotic relative efficiency of the Proschan-Pyke test relative to the Weibull likelihood ratio test when the true distribution is of the form $G(x) = 1 - e^{-(x)^\alpha}$ ($\alpha > 1$) for $x \geq 0$ was computed to be .59 (see Proschan-Pyke (1965)). The asymptotic relative efficiency of their test relative to the gamma likelihood

ratio test when the true distribution is the gamma is only .20. Unfortunately, asymptotic evaluation of the IFR likelihood ratio test seems to be extremely difficult.

There are many additional unbiased tests of exponentiality versus IFRA or IFR which should perhaps be considered. Recall that all of the associated statistics are necessarily homogeneous. A statistic related to the IFR likelihood ratio statistic is

$$\Lambda_n^{***}(\underline{X}) = \left(\frac{n}{\sum_{i=1}^n X_i} \right) \prod_{i=1}^n (n - i + 1) (X_i - X_{i-1}) .$$

If there are no reversals of the normalized differences (they should decrease under IFR alternatives) then Λ_n^{**} and Λ_n^{***} agree except for the factor $n X_1$ and a constant. If $G^{-1}F$ is convex, then

$$\Lambda_n^{***}(\underline{X}) \leq_{st} \Lambda_n^{***}(\underline{Y}) .$$

The test which rejects exponentiality when $\Lambda_n^{***}(\underline{X})$ is sufficiently large is related to a test derived by Moran (1951) for a problem concerning renewal processes. Under the assumption of exponentiality

$$W = \frac{-2 \log \Lambda_n^{***}(\underline{Y})}{1 + \frac{n+1}{6n}}$$

is asymptotically distributed as a χ^2 variable with $n - 1$ degrees of freedom. Epstein's (1960) test 8 uses this statistic. Monte Carlo experiments by Zelen (1961) indicate that the power of this test is less than that of the Proschan-Pyke test for small samples against Weibull distribution alternatives.

In section 2 we proved that $F \leq G$ implies

$$\Lambda_n(\underline{X}) \underset{\text{st}}{\geq} \Lambda_n(\underline{Y}) .$$

Hence we could consider the test, ψ^{**} , which rejects exponentiality in favor of the IFRA hypothesis when $\Lambda_n(\underline{X}) \geq c_{1-\alpha}$ where

$$P_G \{ \Lambda_n(\underline{Y}) \geq c_{1-\alpha} \} = 1 - P_G \{ \Lambda_n(\underline{Y}) \leq c_{1-\alpha} \} = \alpha .$$

For this test we would use the upper percentile points of $-\log \Lambda_n(\underline{X})$ given in Table 1. Since $\Lambda_n(\underline{X})$ is essentially the maximum likelihood under the IFRA assumption it is perhaps not too surprising that it seems superior to the IFR likelihood ratio test (see Fig. 1). On the basis of computer calculations we conjecture that $-\log \Lambda_n(\underline{Y})$, suitably normalized, is asymptotically $N(0, 1)$.

Perhaps a better test than all of those considered so far is a uniform conditional test [see Cox and Lewis (1966) p. 153] based on the mean of the rectangular distribution. This has been described by Bartholomew as the oldest known statistical test [see discussion in Cox (1955)]. Epstein (1960) adapted this test to the life testing problem and called it test 3. The test is based on the total time on test up to the i -th order statistic ($i = 1, 2, \dots, n$), i.e.,

$$T(X_i) = \sum_{j=1}^i (n - j + 1) (X_j - X_{j-1}) .$$

The test statistic is $\sum_{i=1}^{n-1} T(X_i) / \sum_{i=1}^n X_i$. Under the exponential hypothesis

$$Z = \frac{\sum_{i=1}^{n-1} T(X_i) - \frac{(n-1)}{2} \sum_{i=1}^n X_i}{\sum_{i=1}^n X_i \sqrt{(n-1)/12}}$$

is approximately $N(0, 1)$ even for relatively small n . If $F \leq G$, then it follows from Theorem 3.12 (iii) [Barlow and Proschan (1966)] that

$$\frac{\frac{\sum_{i=1}^{n-1} T(X_i)}{\sum_{i=1}^n X_i}}{\text{st}} \geq \frac{\frac{\sum_{i=1}^{n-1} T(Y_i)}{\sum_{i=1}^n Y_i}}{\text{st}} .$$

Hence a natural test, φ^{stat} , rejects exponentiality in

favor of IFRA if

$$\sum_{i=1}^{n-1} \frac{T(X_i)}{\sum_{i=1}^n X_i} \geq c_\alpha$$

where

$$P_G \left\{ \sum_{i=1}^{n-1} T(Y_i) / \sum_{i=1}^n Y_i \geq c_\alpha \right\} = \alpha .$$

Empirical sampling by Zelen and Dannemiller [(1961), p. 47] indicates that this test is superior to φ^{stat} against Weibull distribution alternatives. Investigations by Cox (1955) show that the analogue of this test for randomness in a sequence of events is the most powerful test of the Poisson hypothesis against the alternative of a time-dependent Poisson process with occurrence rate

$$\lambda(t) = e^{\alpha + \beta t} .$$

See Bartholomew (1956) for further results concerning this test.

7. Concluding Remarks

It is perhaps worth noting that the percentage points in Table 2 and the results of section 5 also apply to the Boswell test for trend in a stochastic process of Poisson type. However, if the sample size is n and one is using the Boswell statistic then one should locate percentage points in Table 2 corresponding to

The maximum likelihood estimate assuming a decreasing density is given by Grenander (1956). Recall that if F is DFR, then it has a decreasing density.

Likelihood ratio tests for the two sample problem will be considered in a subsequent paper.

20,000 Random Simulations
for Each Point

$n = 10$




Fig. 1. Power comparison of IFR likelihood ratio test, Proschan-Pyke test and ϕ test against Weibull distribution alternatives. (α is the Weibull shape parameter.)

REFERENCES

- [1] Ayer, M., Brunk, H. D., Ewing, G. M., Reid, W. T. and Silverman, E. (1955).
An empirical distribution function for sampling with incomplete information.
Ann. Math. Statist. 26 641-647. ~~~
- [2] Barlow, R. E. and Proschan, F. (1966). Inequalities for linear combinations of
order statistics from restricted families. Ann Math. Statist. 37 1574-1592. ~~~
- [3] Barlow, R. E. and Proschan, F. (1965). Mathematical Theory of Reliability.
J. Wiley & Sons, New York.
- [4] Birnbaum, Z. W., Esary J. and Marshall, A. W. (1966). Stochastic characteri-
zation of wear-out for components and systems. Ann Math. Statist. 37
816-825. ~~~
- [5] Boswell, M. T. (1966). Estimating and testing trend in a stochastic process of
Poisson type. Ann. Math. Statist. 37 1564-1573. ~~~
- [6] Grenander, Ulf. (1956). On the theory of mortality measurement. II. Skand.
Aktuarietidskr. 39 125-153. ~~~
- [7] Kendall, M. G. (1938). A new measure of rank correlation. Biometrika. 30
81-93. ~~~
- [8] Kiefer, J. and Wolfowitz, J. (1956). Consistency of the maximum likelihood
estimator in the presence of infinitely many incidental parameters.
Ann. Math. Statist. 27 887-906. ~~~
- [9] Mann, H. B. (1945). Nonparametric tests against trend. Econometrica. 13
245-259. ~~~
- [10] Marshall, A. W. and Proschan, F. (1965). Maximum likelihood estimation for
distributions with monotone failure rate. Ann Math. Statist. 36 69-77. ~~~
- [11] Marshall, A. W., Walkup, D. W. and Wets, R. J. B. (1966). Order-preserving
functions; applications to majorization and order statistics. Boeing
Scientific Research Document D1-82-0583.
- [12] Neyman, J. and Pearson, E. S. (1928). On the use and interpretation of certain
test criteria for purposes of statistical inference. Biometrika. 20, Pt. I
175-240, Pt. II 263-294. ~~~
- [13] Neyman, J. and Pearson, E. S. (1933). On the problem of the most efficient
tests of statistical hypotheses. Trans. Roy. Soc. London. Series A,
231 289-337. ~~~

REFERENCES

- [14] Ostrowski, A. (1952). Sur quelques applications des fonctions convexes et concave au sens de I. Schur. J. Math. Pure Appl. 31 253-292.
- [15] Proschan, F. and Pyke, R. (1956). Tests for monotone failure rate. Proc. 5th Berkeley Symp. Math. Statist. and Prob. (to appear).
- [16] Wilks, S. (1962). Mathematical Statistics, J. Wiley & Sons.
- [17] Marshall, A. W. and Proschan, F. (1967). Maximum likelihood estimation for distributions with increasing or decreasing failure rate average. (To appear as a Boeing Scientific Research Laboratories Document.)
- [18] Barlow, R. E. and Jacobson, L. Robustness study of distribution-free life test procedures. (To appear.)
- [19] Moran, P. A. P. (1951). The random division of an interval. Pt. II. J. R. Statist. Soc. B 13 147-150.
- [20] Epstein, B. (1960). Tests for the validity of the assumption that the underlying distribution of life is exponential. Pts. I and II. Technometrics 2 83-101 and 167-183.
- [21] Zelen, M. and Dannemiller, M. C. (1961). The robustness of life testing procedures derived from the exponential distribution. Technometrics 3 29-49.
- [22] Cox, D. R. and Lewis, P. A. (1966). The Statistical Analysis of Series of Events. Methuen Monograph, J. Wiley & Sons.
- [23] Cox, D. R. (1955). Some statistical methods connected with series of events. J. R. Statist. Soc. B 17 129-164.
- [24] Bartholomew, D. J. (1956). Tests for randomness in a series of events when the alternative is a trend. J. R. Statist. Soc. B 18 234-239.

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R&D		
<small>(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)</small>		
1 ORIGINATING ACTIVITY (Corporate author) University of California		2a REPORT SECURITY CLASSIFICATION Unclassified
		2b GROUP
3 REPORT TITLE LIKELIHOOD RATIO TESTS FOR RESTRICTED FAMILIES		
4 DESCRIPTIVE NOTES (Type of report and inclusive dates) Research Report April 1967		
5 AUTHOR(S) (Last name, first name, initial) Barlow, Richard, E.		
6 REPORT DATE April 1967	7a TOTAL NO OF PAGES 29	7b NO OF REFS 24
8a CONTRACT OR GRANT NO Nonr-3656(18)	9a ORIGINATOR'S REPORT NUMBER(S) ORC 67-16	
b PROJECT NO NR 042 238	9b OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
c		
d		
10 AVAILABILITY LIMITATION NOTICES Distribution of this document is unlimited		
11 SUPPLEMENTARY NOTES		12 SPONSORING MILITARY ACTIVITY Office of Naval Research Mathematical Science Division
13 ABSTRACT Likelihood ratio tests for the problem: $H_0: F \in \mathcal{F}_1$ versus $H_1: F \in \mathcal{F}_2$ are defined for certain nonparametric families of distributions and \mathcal{F}_1 . In particular the likelihood ratio test is defined and shown to be unbiased when \mathcal{F}_1 denotes the exponential distributions (possibly truncated) and \mathcal{F}_2 denotes the distributions with increasing failure rate. Comparisons are made with competing tests. The problem of testing for increasing failure rate average is also examined.		

DD FORM 1473

Unclassified

Security Classification

14 KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Likelihood ratio tests increasing failure rate life testing nonparametric statistics						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (either by the originator or by the sponsor), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.